

1. Assess airway and initiate basic and/or advanced airway maneuvers prn (*MCG 1302*)
2. Prioritize treatment of systemic symptoms  
For signs or symptoms of allergic reaction, treat in conjunction with *TP 1219, Allergy*  
For poor perfusion, treat in conjunction with *TP 1207, Shock/Hypotension*
3. Keep patient calm and limit activity  
Position affected extremity at or below level of the heart
4. For SNAKE bites:  
Splint the affected area  
Elevate the extremity to the level of the heart
5. For INSECT (bee, wasp, ant), spider and scorpion stings:  
Remove stinger if visualized ❶  
Apply cold pack
6. For MARINE envenomation (e.g., jelly fish, stingrays and scorpion fish):  
Remove barb when applicable  
Soak area in hot water if available ❷
7. Establish vascular access prn (*MCG 1375*)
8. For continued pain after specific measures above: (*MCG 1345*)  
**Fentanyl 50mcg (1mL) slow IV push or IM/IN**  
Repeat every 5 min prn, maximum total dose prior to Base contact 150mcg  
**Morphine 4mg (1mL) slow IV push or IM**  
Repeat every 5 min prn, maximum total dose prior to Base contact 12mg  
  
**CONTACT BASE** for additional pain management after maximum dose administered:  
May repeat as above up to maximum total dose Fentanyl 250mcg or Morphine 20mg
9. For nausea or vomiting:  
**Ondansetron 4mg ODT/IV/IM**, may repeat x1 in 15 min prn

**SPECIAL CONSIDERATIONS**

- ❶ Remove stinger by scraping patient's skin with the edge of a flat surface (credit card or similar). Do not attempt to pull the stinger out with fingernails or tweezers, as this may cause release of additional venom.
- ❷ Do not use vinegar given the type of jellyfish endemic to California.